



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

### A New *Eleocharis*.

By CHARLES WRIGHT.

*ELEOCHARIS DIANDRA*.—Culms nearly terete (8'-10' high) from tufted, fibrous roots, rather slender; spike ovoid, very obtuse (2"-4" long); scales numerous (80-120) ovate, rounded at the tip, one-third longer than the achenium, imbricated in many rows; stamens two; achenium cuneate-obovate, pale stramineous becoming castaneous, shining, scarcely half a line long; bristles few and short, or mostly wanting; tubercle depressed, transversely oblong, the ends rounded and slightly elevated, the top shortly apiculate; style usually bifid.

On high sand-bars of the Connecticut River, between Hartford and Wethersfield, growing in company with *E. obtusa*, which it closely resembles in appearance. A tabular statement of the differences is appended.

	<i>Eleocharis obtusa</i> , Shultes.	<i>Eleocharis diandra</i> , n. sp.
TUFTS mostly	larger, annual.	smaller, annual.
SPIKE	very blunt.	bluntish or acutish.
SCALES	rounded at top; one-third longer than the achenium.	narrowed towards the blunt apex; twice as long as the achenium.
STAMENS	three.	two.
BRISTLES	exceeding the tubercle.	few and short or mostly O.
ACHENIUM	larger, tapering evenly to the base.	smaller, tapering more abruptly into a narrower base.
TUBERCLE	crest-like and thin, curved on the lower edge, which rests its whole length on the achenium, and is almost as broad, the two other sides forming about a right angle.	transversely oblong, its end rounded and raised above the achenium, and little more than half as broad, shortly apiculate.
STYLE	3-cleft, rarely bifid.	bifid, rarely 3-cleft.

The main differences are: first, *two* stamens, for which in our floras there is no generic allowance; secondly, the usually bifid style; thirdly, the tubercle wanting the *cusps* which run down the top of the achenium of *E. obtusa*; and, fourthly, the few small bristles, or mostly none at all.

### New or Little-known Ferns of the United States, No. 14.

By D. C. EATON.

53. *Phlegopteris reptans*, (*Polypodium reptans*, Swz.; *Aspidium reptans*, Mett.),—Rootstock short, creeping; stalks clustered, gray-stramineous, slender, naked, a few inches to a foot long; frond as long as the stalk, membranaceous, softly hairy with branched or stellate hairs, oblong lanceolate, pinnate with nearly or quite sessile oblong or sometimes rounded obtuse crenately pinnatifid pinnæ, the apex pinnatifid and often elongated and rooting; veins pinnate, simple, the basal veinlets often anastomosing; sori rather small, seated on the middle of the veinlets, naked or with a minute rudiment of an indusium.